PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS — CAUTION: Harmful if swallowed, absorbed through skin or inhaled. Avoid contact with er after handling and before eating, drinking, chewing gum, using tobacco, or using the

PERSONAL PROTECTIVE EQUIPMENT (PPE):

All mixer/loaders and applicators must wear a minimum of a NOSH-approved particulate filtering facepiece respirator with any N, R or P filte OR a NOSH-approved elastometric particulate respirator with any N, R or P filter, OR a NOSH-approved power dair-purifying respirator with a 1-F filter. Repeated reposure to high concentrations of microbil proteirs can cause allergic sensitazion. Follow the manufacturer's instruction. for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately

ENGINEERING CONTROLS: When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Protection Standard (WPS) for agricultural pesticides (40 CFR 170.607(d) and (e)), the handler PPE requirements may be reduced or

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

USER SAFETY RECOMMENDATIONS

Remove dothing/PPE immediately if pesticide gost inside. Then wash throroughly and put on clear usuring.
 Remove PPE immediately after trading this product. Wash the outside of gloves before removing. As soon as possible, wash throroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS: For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or this, cross title state or thick agency reprovises for prescribed progulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard controllural profession of aminuflural workers on farms, forests pursavies, and properly uses and bandlers of agricultural workers on farms. requestions our empression of adjustation state of the st

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has

- Coverails
 Waterproof gloves
 Shoes plus socks

PRODUCT INFORMATION

- LIASTOP CONTANS W6 is a water dispensible granule for the reduction/control of Scientinia scientiniam and Scientiniam innor in agricultural, nursery and greenhouse soils.

 LIASTOP CONTANS W6 is registered to be used on soils where plants susceptible to Scientinia scientiniam and Scientiniam innor are grown. These plants include trassist, colleit leafly registables, but vegetables, cered grains, cucurativ registables, fluting vegetables, hefts and spices, leafly vegetables, legure vegetables, sonor-grass animal feeds, obsect crops, ornamentals, root and tuber vegetables. For use in pre-plant, in crop, post-harvest by toroadcast, and band, See April 2012. The NTO-MEDIATION NFO-MEDIATION NFO-MEDI

MIXING DIRECTIONS

Determine the volume of water needed to provide thorough coverage of the soil in the treatment area. The amount of water needed will depend on the weather, spray equipment and local experience. Partially fill the spray tank with clean water and begin agilation. Add the specified amount of LASIGPO CONTRON Will to the text (covoral application decircions below). Final filling the text for the sciented volume that provides maximum coverage. Maintain agitation throughout spray application. Do not allow spray mixture to stand overnight or for prolonged periods.

APPLICATION INFORMATION

- APPLICATION INFORMATION

 LASTOP CONTANS Wis dispersed in water and applied directly to the soil surface, using conventional spray equipment.

 Incorporate LASTOP CONTANS Wis soon after application into the upper 1 to 2 inches of soil by means of studior mechanical incorporation by disc. cultivator, lifer or other similar equipment who available, or by overhead irrigation where available, or rain water. For specific application situations and environments, does us applications options with your field advisor or product distributor.

 When applied correctly, LASTOP CONTANS Wis will reduce incodum back of Scientifia activated and hybrise in the treated soil and, if water breast, or one probability, and soil reduces a product entering the soil.

 Apply LASTOP CONTANS Wis you elicited surger or dents to the sol prior to planting, or at planting or after harvest on crop studble.

 Apply LASTOP CONTANS Wis give a licetaced surger or dents to the sol prior to planting, or at planting or after harvest on crop studble.

 Apply LASTOP CONTANS Wis give increase elicited to to todain optimal efficacy.

 Use higher application rates when the weather conditions are expected to be conductive for disease development, if the size of the soil surface, and in minure is present, or if minimum/bowfor-lill programs are endined.

 Apply LASTOP CONTANS Wis Girectly to the soil surface and incorporate into the size of the programs are endined into the upper 1 to a riches of soil soon after application, when possible. The program is programs and the programs are endined in the programs are endined in the program of the programs are endined in the program of the programs are endined in the programs. The programs are endined in the programs. The programs are endined in the programs are endined in the programs are endined in the programs. The programs are endined in the programs ar

- dissipated.

 ALSTOP CONTANS WG has no qurative effect and therefore is not effective appirist plants infected with disease at the time of application.
 For maximum effectiveness, follow up an initial pre-planting treatment with a treatment of the corp shubble after harvest of the Scienotiniasusceptible crop, Repeated use throughout the rotation increases long-term efficacy due to reduction of the pathogen.
 Apply the higher rate when known Scienotinia mixor exists in the soil.

 For ground towardset applications, apply LALSTOP CONTANS WG as a spray at a minimum volume of 10 galons of water per acre prior to, at planting, after transpart, or post-travers. Maintain adjustion or hing spray operation.

 Through and uniform coverage of the soil surface is necessary for control of S. sclerotorum as surviving sclerotal produce accesspores which become alrohome and stored nor the entire surface.

- indrough and uniform coverage or the soft startes is necessary for corror of a Scienterorum's surviving sciential produce ascuspores which become artistome and spread over the entire surface. When applying LASTOP CONTAKS WG on crop stubble of susceptible crops and prior to replant of a susceptible crop, use a rate of 1 to 20 bit of product per acre paraglication to soll after crop harvest. Allow adequate growing season time (iminimum 2 to 3 months) and/or soil temperature for LASTOP CONTAKS WG to remain actively growing to colonize and infect. Scientific scientials Apply LASTOP CONTAKS WG to minimum-fill or no-fill fields, or on crop stubble. LASTOP CONTAKS WG efficacy will be increased when application is done prior to rain or any stubble treatment, as this will increase distribution and reach of the spores.

- Do not apply LALSTOP CONTANS WG via broadcast or sprinkler at time of plant emergence until harvest. Only use LALSTOP CONTANS
 WG by sol-directed drip or trickle irrigation methods at time of plant emergence until harvest.
 Do not tark- mix with, or apply LALSTOP CONTANS WG within 7 days before or after use of other fungicide products.
 For information on which pescidices can be mixed with LASTOP CONTANS WG without harming the beneficial fungus it contains, contact
 your LALLEMAND Technical Sales Representative [or Spoam Agro USA, Inc.].

For agricultural uses - Apply LALSTOP CONTANS WG to the soil at a rate of 1 to 4 to per acre per application prior to planting, or at planting or after transplant, or postranest. LALSTOP CONTANS WG may also be applied to plant debris that remains in the field after harvest, prior to replart of a susceptible crop, if incorporation will displace the soll greater than two inches, increase the application rate to 3 to 6 to, per acre. Make a maximum of eight (8) applications of LASTOP CONTANS WG per growing season or per year, at labeled rates, as required to maintain disease control. This product may be applied up to the day of harvest on all crops.

For greenhouse, covered crops, and nursery uses - Apply LALSTOP CONTANS WG at a rate of 0.75-1.5 oz. per 1000 sq. ft. Make a maximum of eight (8) applications of LALSTOP CONTANS WG per growing season or per year, at labeled rates, as required to maintain disease control. This product may be applied up to the day of harvest on all crops.

Susceptible Plants Brassica (Cole) Leafy Vegetables: Broccoli, Chinese broccoli (gai lon), broccoli raab (rapini), Brussels sprouts; cabbage; Chinese cabbage (bok choy, napa), Chinese mustard cabbage (gai choy), cauliflower, cavalo broccolo, collar mustard greens, mustard spinach, rape greens; including cultivars, varieties, and/or hybrids of these

Bulb Vegetables: Chive, fresh leaves; chive, Chinese, fresh leaves; daylily, bulb; elegans hosta; fritilaria, bulb; fritilaria, bulb; fritilaria, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek, wild; lily, bulb; onion, Beltsville bunching; onion, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; kurrat; lady's leek; leek; leek; wru; iny; uou; union; uenovine downing, downing, bulb; orition, Chirese, bulb; orition, fresh; orition, green; orition, macrostem; orition, pearl; orition, potato, bulb; orition, fresh kavies; outlivars; varieties, and/or hybrids of frese

Cucurbit Vegetables: Chayote (futil), Chinese waxgourd (Chinese preserving melon), citron melon, oucumber, gherkin, edible gourd (includes hydran, cucurza, herbinna, Chinese okra), Momardica spp (includes balsam apple, balsam pear, billstimelon, Chinese oucumber), muskmelon (includes the caraldoupe, carladoupe, casabla, Santa Caius melon, cerishaw melon, honeyslaw melon, troneyslab. Persiam melon, golden persiawa melon, mango melon, priseaple melon, raske melon, and other varieties and/or hybrids of these), pumpkin, squash (summer, winter which includes butternut squash, calabaza, hubbard squash, acom squash, spaghetti squash), watermelon; cultivars, varieties, and/or hybrids of these

Fruiting Vegetables (except cucurbits): Eggplant, groundcherry (Physalis spp), pepino, pepper (includes bell pepper, chili pepper, chiling vegetables (except cucurbits): Eggplant, groundcherry (Physalis spp), pepino, pepper (includes bell pepper, chiling pepper):

Herbs and Spices: Allspice, angelica, anise (including star), annatto (seed), balm, basil, borage, burnet, chamomile, caper buds; Herbs and Spices: Alspice, anglaica, arises (including star), amatilo (seed), baim, bask to broage, burnet, charomatic caper busk caratherom, cassis bark; cassis bunks; cathio; celary seed; chevul (fried); chive, chive, chives; cinnamo clary, chove busk; coraindar leaf cilcantro or Ohinese parasely; coraindar seed cilcantro); costamary, culamto (sead); culamto (sead); culamto (sead); chiven (common, term); celar, di (dilveved; di (sead); fremel (common); fremel Puronos (seed); freuquesk; quaris or parasitis, horehound; hysson; juriper benry, tevende; kemongrass; lovage (sead); lovage (sead); mace; marigoid, marjoram; mustard (sead); nashufrum; nutmeg, paraley (dried); pennyroyal; pepper, black; pepper, white; poppy (sead); rosemary; rue; saffron; sage; savory, summer an writer; sweet losy; tasny; cultivas; varieties; and for hybrids of these

Leafy Vegetables (except brassica vegetables): Amaranth (Chinese spinach), augula (roquette), cardoon, celery (including Chinese), calubor, cherul, chrysamtherum (ecitibe-leaved, garland), com salad, cress (garlan, upland), dardellon, dock (sorrel), endive (escardie), Florence Fernel, lettuce (head, leaf), orach, pastey, pursiane (garden, winter), radicchio (red chicory), rhubarb, spinach (including New Zelland, vine), Swiss chard, cultivars, varieties, and/or hybrids of these

Foliage of Legume Vegetables: Any cultivar of bean (Phaseolus), field pea (Pisum) and soybean. Plant parts of any legume e legume vegetables that will be used as animal feed; cultivars, varieties, and/or hybrids of the

Legume Vegetables (succulent and dried): Bean (Lupinus) (includes grain, sweet, white, and white sweet), bean (Phaseolus) Leguinir vegetaties (socional and unles) pear (capital) quotous graft, siever initie, and wite siever), learl (viscous) (includes field), which, yim, raw), printo, runner, stap, tepany, wan), bean (Vigra) (includes adauk, asparajus, biokeyed pea, calif, Chinese (onghean, cowpea, cowider pea, moth, mung rice, southern pea, und, yardrong), broad bean (fana), chickpea (garbanz quar, adadean, blabb bean, lent), pea (Fesm) (includes whart deliber, podied, regiller), field; pear (regen, stow, sugar strap), pigeon pea, soybean, soybean (immature seed), sword bean; cultivars, varieties, and/or hybrids of these

Non-grass Animal Feeds (forage, fodder, straw, hay): Alfalfa, velvet bean, clover (*Trifolium*, Melliotus), kudzu, lespedeza, lupin, sainfoin, trefoil, vetch (including crown, milk); cultivars, varieties, and/or hybrids of these

Oilseed Crops: Borage, calendula, castor oliplant, Chinese tallowiree, cottonseed, crambe, cuphea, exhium, euphorbia primrose, likar seed, gold of pleasure, hare's ear mustant, jojoba, lexpuereila, lunaria, meadowloam, milkweed, mustant seed, oil radish, poppy seed, rapesseed, rose hip, safflower, sesame, stokes aster, sunflower, sweet rocket, tallowwood vernonia; cultivas, vatellea, and/or hiphids of these

Ornamental Herbaceous Potted Flowers and Bedding Plants: Chrysanthemums, cy petunia, poinsettia, primrose, snapdragon; cultivars, varieties, and/or hybrids of these

cassava, bitter and sweet; celenac; chervil, turnip-rooted; chicory; dashe salsify, black; sweet potato; tanier; turnip; yam, true; cultivars, varieties, ar

light mechanical incorporation or by irrigation application, especially in warm, dry climates.

Application Instructions

Use Rate/Application

(field and chemigation applications)

0.75 to 1.5 oz/1000 ft²

applications)

Chemigation applications: This covers all methods of rrigation described as allowed in the text of this label

For field and chemigation applications: apply the higher For near and consimpagnic approaching a plant and a rate when Scientific more is present in field. Apply minimum 1 to 4 lbA 7 days prior to planting or at planting, or soon after planting followed by 1 to 4 lbA 1 to 21 days after first spray, or just after thinning, Incorporate into top 1 to 2 inches of soil Immediately after each application when possible. On ord train-risk with other fungioides fostation with other fungioides allowed after 3 weeks following an application of this product.

In Leafy Vegetables (except brassica vegetables): apply In Leasy vegetanes experient indicates and a second application at least 2 IbA immediately after trinning followed by a second application 14 days later. Make broadcast or banded application with a minimum of a 12 total band with trageled at the crop stem. Incorporate LALSTOP CONTANS WG into top 2 inches of soil by joyth mechanical incorporation or by imgation or rainfall soon after application, especially in weim, dry climates. Fields in which Scientifia minor's present, apply at least the 2.0 bl/x rate or hother in-field or chemination anotherisers inmediately and ninning followed by and application to plant debris immediately

Apply spray or drench to plug trays or pots. Apply the hi when Soleratinia milror is expected after transplanted Water in immediately after each application. Do not tank other fungicides. Rotation with other fungicides is allowe 3 weeks following an application of this product.

Incorporate into top 2 inches of soil immediately after each application when possible. Do not tank-mix with other fungicides Rotation with other funcicides is allowed after 3 weeks following an application of this product.

Chemigation applications: This covers all methods of irrigation described as allowed in the text of this label.

For field and chemigation applications: apply the higher rate when Sclerotinis minor's gresent in field Apply minimum I ad 640.7 dep prior biganting or at planting, or somo after planting followed by 1 to 4 tb/A 14 to 21 days after first stray, or just after thirning, incorporate into for 1 to 2 inches soil immediately after each application when possible. Do not tark-rinx with other funglocks is of allowed after 3 weeks following an application of this product.

other fungicides. Rotation with other fungicides is 3 weeks following an application of this product.

CHEMIGATION INSTRUCTION

stems either pre-plant or to crop stubble on lighter textured soils (sandy loams – sand) at a rate appropriate to th 34 –1 inch of water will be enough for a good incorporation of LALSTOP CONTANS WG. If followed by tillage, do

Mixing instructions: A ANS WG in the supply tank to a concentration appropriate to cover the intended acreage in one pass (example: 40 acres at 2 lb. per acre will require 80 pounds spension of the spores of the active agent in the injection solution. All of the LALSTOP CONTANS WG should be applied in one pass of the irrigation system.

r including center pivot, lateral move, side (wheel) roll irrigation systems. Do not apply this product through any other type of irrigation system.

rop can result from non-uniform distribution of treated water. stem, you should contact your State Extension Service specialists, equipment manufacturers or other experts.

responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

m for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days

ad to public water systems must contain a functional, reduced pressure zone, back-flow preventer (RPZ) or the functional equivalent in the water supply line upstream from the public water system should be discharged into a reservoir tarik prior to posticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of tark of at least twice the inside diameter of the fill pipe.

pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the intigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point w

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system inte

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry cool place out of direct sunlight and away from heat sources at approximately 39 °F. Keep from overheating. (Almacene en un lugar fresco seco fuera de luz directa del sol y lejos de fuentes de calor en aproximadamente 39 °F. Mantenga de recalentar.)

CONTAINER HANDLING: Non-refilable container. Do not reuse or refill this container.

Bag: Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke Plastic container: Clean container promptly after emptying. Tirple-rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds Pour insate into application equipment or a mix tank or store insate for later use or disposal. Dark for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available, or dispose of empty package in a sanitary and roll, or by inchination, or, if allowed by state and local authorities by, burning if turned, sign of all smoke.

NOTICE TO LISER

Darstar Ferment AG / LALLEMAND PLANT CARE warrants only that this product conforms to the product description on this label and is reasonably lift for the purposes set forth in the Directions for Use when used in accordance with them. However, ineffectiveness or other unintereded consequences may result because of such factors as the use, storage or handling of the product contract by the label instructions, all of which are beyond the control of Denstar Ferment AG / LALLEMAND PLANT CARE shall not be label for indirect or consequentful damages resulting from the use, storage or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, DANSTAR FERMENT AG / LALLEMAND PLANT CARE MAKES NO WARRANTIES OF MEDICHANIZBULTY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE



LALLEMAND

Net Weight: 25 lb (11.35 kg)